



P-ISSN: 2664-3685
E-ISSN: 2664-3693
www.paediatricjournal.com
IJPG 2018; 1(2): 36-39
Received: 09-09-2018
Accepted: 17-10-2018

Dr. Oniell Fernandes
Assistant Professor,
Department of Paediatrics,
Kanachur Institute of Medical
Sciences, Mangalore,
Karnataka, India

Study of outcome of meconium stained amniotic fluid in newborn

Dr. Oniell Fernandes

DOI: <https://doi.org/10.33545/26643685.2018.v1.i2a.178>

Abstract

Background: The presence of meconium stained amniotic fluid (MSAF) is a serious sign of fetal compromise, which is associated with an increase in perinatal morbidity, clear amniotic fluid on the other hand is considered reassuring. A study by Mathews and Warshaw found that in 98.4% of cases with meconium stained liquor, the neonates were admitted to the neonatal intensive care units, these were delivered at 37 weeks of gestational age or later as it shows maturation of autonomic nervous system. MSAF occurs in 10% of pregnancies with most occurring at term or particularly post term. The purpose of this study was to evaluate the effect of clear liquor and meconium stained liquor on time and mode of delivery and to find out neonatal morbidity and mortality associated with meconium stained amniotic fluid.

Keywords: Meconium stained liquor, LSCS, Newborn outcome, meconium aspiration syndrome

Introduction

Meconium is named according to Greek literature “mekonionarion” – “opium-like” as said by Aristotle in reference to its tarries like appearance and a belief that it induces sleep in the fetus [1]. The passage of meconium into the amniotic fluid during labor (fresh meconium) is one of the traditional indicators of fetal distress and is associated with increased fetal morbidity and mortality [2]. Meconiumstained amniotic fluid (MSAF) is known to be associated with several maternal and neonatal risk factors [3]. In healthy, well-oxygenated fetuses, meconium is cleared from the lungs by the normal physiological mechanism. Most of them can develop an effective breathing movement to signs of aspiration and persistent hypoxia [4]. MSAF is considered a harbinger of fetal compromise because of its direct correlation with fetal distress and increased likelihood of aspiration of meconium with resultant deleterious effects on the neonatal lung [5]. Meconium passage is rare before 34 weeks of gestations and after 37 weeks its incidence increases steadily with increasing gestational age [6]. Passage of meconium in utero with staining of amniotic fluid occurs in 12% to 16% of all deliveries [7-10]. Presence of meconium below the vocal cord is known as meconium aspiration and this finding occurs in 20% to 30% of all infants with MSAF fluid with around 12% mortality [8,10].

The purpose of this study was to evaluate the effect of clear liquor and meconium stained liquor on time and mode of delivery and to find out neonatal morbidity and mortality associated with meconium stained amniotic fluid.

The purpose of this study was to evaluate factor affecting incidence of MSAF and to find out neonatal morbidity and mortality associated with meconium stained amniotic fluid.

Materials and methods

Descriptive observational study was conducted in the pediatric Department at tertiary care hospital over a period of one year (August 2016 to August 2017). A total number of meconium stained amniotic fluid (MSAF) were included in the study out of 10800 live birth.

Inclusion and Exclusion criteria: Inclusion criteria included all the newborn with meconium stained amniotic fluid (MSAF) delivered in our tertiary care hospital.

Exclusion criteria were all the newborn with meconium stained amniotic fluid (MSAF) delivered outside our tertiary care hospital and admitted in NICU and those with major congenital anomaly.

Corresponding Author:
Dr. Oniell Fernandes
Assistant Professor,
Department of Paediatrics,
Kanachur Institute of Medical
Sciences, Mangalore,
Karnataka, India

Methods

A total number of 940 babies born of Meconium stained amniotic fluid(MSAF) were included in the study and those admitted to NICU were studied in a detailed manner; history was noted, clinical examination was done; they were investigated as per proforma and followed up till the time of discharge. Also included newborn who are presented with respiratory distress soon after birth and/or x-ray chest showing evidence of aspiration pneumonitis (atelectasis/hyperinflation) are considered as Meconium Aspiration Syndrome (MAS).

All patients during the study were managed according to neonatal resuscitation guideline 2017, babies with MSAF divided into vigorous and non-vigorous groups and managed accordingly [1].

In each and every patient with history of meconium stained amniotic fluid, APGAR was counted at 1 min & 5 min, if 5 min APGAR was < 7 then APGAR at 10 min & 20 min counted. Requiring positive pressure ventilation and suction was solely on the basis of status of the baby at the time of birth, whether vigorous or non-vigorous and then observed for development of signs and symptoms of respiratory distress and graded as per downe’s scoring system as mentioned in the table. X-ray chest after 6 Hrs of birth in each cases with increased respiratory rate.

Sarnet and Sarnet staging for hypoxic ischemic encephalopathy

Feature	Stage 1	Stage 2	Stage 3
Consciousness	hyperalert	lethargic	Comatose
Muscle tone	normal	lethargic	Flaccid
Deep tendon reflexes	Brisk	exaggerated	Absent
Myoclonus	present	present	Absent
Sucking	normal	weak	Absent
Moro response	exaggerated	incomplete	Absent
Oculocephalic reflex	normal	Over reactive	Reduced/absent

We also observed for development of other complication like pneumothorax, septicaemia, metabolic acidosis and severe PPHN

Result

A total of 940 babies born with history of MSAF and admitted to our hospital were studied. Incidence of MSAF is 8.7% in present study [7].

Total Delivery during the Study = 10800

Total New-born with MSAF = 940

Total NICU Admission = 4308

Total MSAF Requiring NICU Admission = 560 (13% of Total Admission)

Incidence of MSAF in Present Study = 8.7 %

Table 1: Comparison of gender prevalence

	MSAF	Total Delivery
Male	520 (9.03%)	5758
Female	420 (8.33%)	5042
Total	940	10800

Using chi-square test for statistical significance, p value is 0.25 which suggest that there is no significant correlation between incidence of MSAF in male and female.

Downe’s scoring system

Score	0	1	2
Respiratory rate/min	< 60	60-80	>80
Cyanosis	None	In room air	In 40% fio2
Retraction	None	mild	Severe
Grunting	None	Audible with stethoscope	Audible without stethoscope
Air entry	Clear	Bilaterally decreased	Barely audible

Score

1-3: mild respiratory distress

4-7: moderate respiratory distress- prepare to transfer/intubate baby

>7: impending respiratory failure- intubate

Babies without respiratory distress and who were vigorous at birth were shifted to mother side, breast feeding established and immunized with OPV, BCG & Hepatitis-B Vaccine.

All babies who required some form of resuscitation or who developed signs of respiratory distress were shifted to NICU and managed accordingly. Babies were observed for development of hypoxic ischemic encephalopathy as defined by Volpe. Those who developed HIE were scored according to Sarnet & Sarnet scoring system as per table.

Table 2: Comparision of gestation

Maturity	MSAF	Percentage
Preterm	42	4.5%
Full Term	825	87.8%
Post Term	73	7.7%
Total	940	

According to the “New Ballard Scoring” the babies were divided into the following gestational categories.

In the present study when maturity of the fetus and MSAF are considered together, there were 4.5 % of pre-term, 87.8 % were full term & 7.7 % were post term.

Table 3: Association between mode of delivery

Mode of Delivery	Vaginal	LSCS	Forceps	Vaccum
Normal (9860)	8039(81.5%)	2548(25.8%)	51(0.5%)	162(1.6%)
MSAF (940)	532(56.6%)	395(42.0%)	10(1.1%)	3(0.003%)

When compared between normal vaginal and LSCS, LSCS (especially done in emergency) is done more commonly in the patients with MSAF.

Table 4: Comparison of maternal risk factor (N=940)

PIH	269 (28.6%)
PROM	190(20.2%)
Oligohydroamnios	118(12.5%)
Post Date	73(7.7%)
APH	72(7.7%)
Eclampsia	32(3.4%)
Severe Anemia	8(0.01%)
Elderly Primi	4(0.004%)
No Risk Factor	174(18.5%)
Total	940

Out of 940 babies in the study mothers of 766(81.49%) had one of the risk factors, mentioned in the chart. Of which 28.6% had PIH, 20.2% had PROM, and 12.5% had oligohydroamnios. This correlates with the findings of high risk situations with increased likelihood of in utero passage of meconium passage, which if timely intervened can decrease the incidence of morbidity and mortality among the babies [8].

Table 5: Comparison of birth asphyxia

Apgar At 5 Min.	Developed Mas	Not Developed Mas	MSAF
Severe birth asphyxia(1-3)	9(3.6%)	0	9
Moderate birth asphyxia(4-6)	144(57.6%)	166(24.1%)	310
Mild birth asphyxia(7-10)	97(38.8%)	524(75.9%)	621
Total	250	690	940

In present study out of 940 babies with MSAF 250 (26.6%) developed MAS. When APGAR score at 5 min and development of MAS considered, among 1 to 3 scored group 100% developed MAS, in 4 to 6 scored group 46.5%, in 7 to 10 scored group 15.6% developed MAS [9].

When grouped in to < 7 and 7 or more scored groups, < 7 scored group constituted 153 (61.2%) babies out of 250 babies with MAS. These finding indicates that incidence of MAS is significantly high in babies born through MSAF with APGAR at 5 min < 7, but score of 7 or more does not rule out chance of developing MAS because MAS starts as an intrauterine event.

Table 6: Complication of MSAF

MAS	250(26.6%)
SEPTICEMIA	153(16.3%)
HIE	84(8.9%)
HIE STAGE 1	42
HIE STAGE 2	26
HIE STAGE 3	16
SYMPTOMATIC PPHN	24(2.6%)
PNEUMOTHORAX	9(1%)

From 940 babies with MSAF, 560 (59.6%) babies required NICU admission for active management. In study done by Erum Majid Shaikh *et al.* out of 250 babies with MSAF 165 (66%) required NICU admission. the cause for higher admission in that study may be due to higher incidence of MAS 115 (46%) as compared to our study in which incidence of MAS is 250 (26.6%). Other complication include sepsis (16.3%), HIE (8.9%) and symptomatic PPHN (2.6%). And 9 patients developed pneumothorax (1%) [8].

Table 7: Oxygen requirement

Oxygen Requirement	MSAF newborn
CPAP	564(60%)
Ventilatory Care	188(20%)
None	188(20%)
Total	940

Table 8: Outcome of ventilated patients

Survived	Expired	Total
26(13.8%)	162(86.2%)	188

In the present study out of 940 babies, total 188(20%) babies requires ventilator support due to complication of

MSAF; Of all babies who required ventilator support 26 (13.8%) babies survived and remaining 162 babies (86.2%) expired.

Out of 940 babies 188 babies (20%) does not required any active intervention.

Table 9: Cause of death in MSAF

Cause of death	No. of patients
Septicemia	122
DIC	102
MAS	74
HIE	18
PPHN	15
Pneumothorax	8

Out of 940 babies with MSAF, 146(15.5%) babies expired. Most common cause of death was sepsis 122 patients (13%), followed by DIC 102 patients (10.9%) and MAS 74 patients (7.9%). Other causes include, severe birth asphyxia with HIE (2%), PPHN (1.6%) and pneumothorax (0.09%).

Table 10: Mortality rate in MSAF

Survived	Expired	Total
794(84.5%)	146(15.5%)	940

A total of 146 babies out of 940, died giving an overall mortality rate of 15.5% [10].

Table 11: Mortality rate in MAS

Survived	Expired	Total
176 (70.2%)	74 (29.8%)	250

Out of 250 babies with MAS, 74(29.8%) babies expired.

Discussion

The presence of meconium stained amniotic fluid (MSAF) is a serious sign of fetal compromise, which is associated with an increase in perinatal morbidity, clear amniotic fluid on the other hand is considered reassuring.

In the earlier days, early amniotomy with active management of labour was done to detect meconium passed during labour. Amniotomy in labour is also commonly performed to detect meconium where heart rate is unsatisfactory [11, 12]. If meconium stained amniotic fluid (MSAF) is found, then continuous fetal heart rate

monitoring is required for fetal well being ^[13]. Incidence of MSAF rises with gestational age. A study by Mathews and Warshaw found that in 98.4% of cases with meconium stained liquor, the neonates were admitted to the neonatal intensive care units, these were delivered at 37 weeks of gestational age or later as it shows maturation of autonomic nervous system.

Aspiration of meconium during intrauterine life may result in or contribute to meconium aspiration syndrome (MAS), representing a leading cause of perinatal death ^[14]. MAS is more frequently seen in post term pregnancy or in growth restricted fetuses. Factors such as placental insufficiency, maternal hypertension, pre-eclampsia, oligohydroamnios or maternal drug abuse (tobacco or cocaine) result in, in utero passage of meconium. Meconium stained liquor may be aspirated during delivery resulting in neonatal respiratory distress syndrome ^[15]. The incidence of admission to neonatal intensive care unit with respiratory distress syndrome, meconium aspiration syndrome, neonatal asphyxia, chorio-amnionitis, fetal distress or fetal acidosis were higher in pregnancies complicated by meconium stained liquor.

MSAF occurs in 10% of pregnancies with most occurring at term or particularly post term. Prolonged labour was more common or was associated with worst outcome in meconium stained liquor ^[16]. Caesarean section were performed twice as frequently in women presenting with MSAF and failure to progress, which was the indication in more than half of the cases.

Conclusion

The mortality of babies born with meconium stained amniotic fluid with or without aspiration syndrome is statistically significant which suggest that timely intervention and management can prevent mortality and morbidity.

References

1. Jalal M, Abu Shwaeesh. Diseases of Respiratory system. In: Fanaroff's and martin's Neonatal perinatal Medicine diseases of the fetus and Infant 8th edn. Saunders: Elsevier. 2006;134:563-565.
2. Anne Greenough, Anthony D Milnerpulmonary Diseases of New-born.in: Robert's textbook of neonatology 4th edn. Cruchill living stone. 2006;1199:502-508.
3. Mohamad Reza Sedaghatian, Leena Othman, Nadirea Rashid, Pallath Ramachandran, Abdul Bari Bener. An 8year study of meconium Stained amniotic fluid in different Ethnic groups. Kuwait Medical Journal. 2004;36:266-269.
4. Keenan W. Recommendations for management of the child born through meconium stained amniotic fluid. Pediatrics. 2004;113:133-134.
5. Nizami SQ, Arif MA. Respiratory distress syndrome among hospital born infants. Pak Paediatr J. 1985;9:276-282.
6. Wiswell TA, Bent RC. Meconium staining and the meconium aspiration syndrome, unresolved issues. In pediatric clinics of North America updates on Neonatology. 1993;40:955-981.
7. Brown BL, Gleicher N. Intrauterine meconium aspiration. Obstet Gynecol. 1981;52:26-9.
8. Cochran WG. Sampling techniques. 3rd ed. New York.

John Willey and sons. 177, 81.

9. Orimadegun AE, Akinbami FO, Tongo OO, Okereke JO. Comparison of neonates born outside and inside hospitals in a children emergency unit, southwest of Nigeria. *Pediatr Emerg Care*. 2008;24:354-358.
10. Owa JA, Osinaike AI. Neonatal Morbidity and mortality Nigeria. *Indian J Pediatr*. 1998;65:441-449.
11. American academy of pediatrics, Neonatal Resuscitation 7th edition.
12. Goffinet F, Fraser W, Marcoux S, Bréart G, Moutquin JM, Daris M. Early amniotomy increases the frequency of fetal heart rate abnormalities. *Amniotomy Study Group. Br J Obstet Gynaecol*. 1997 May;104(5):548-53. doi: 10.1111/j.1471-0528.1997.tb11530.x. PMID: 9166195.
13. Miller FC, Sacks DA, Yeh SY, Paul RH, Schiffrin BS, Martin CB Jr, *et al*. Significance of meconium during labor. *Am J Obstet Gynecol*. 1975 Jul 1;122(5):573-80. doi: 10.1016/0002-9378(75)90052-6. PMID: 238396.
14. Cunningham F. Gary: meconijm aspiration. *William obstetrics*, 21st edition (c) chapter 39, 2001.
15. Eichenwald Eric C. meconium aspiration. *Manual of neonatal care* Ed. John p cloherty and Ann R. Stark 6th edition, 2008.
16. Care of baby in the labour room, Meharban singh care of the Newborn, 7th edition.